





A filling rarely chooses a convenient time to fail. Most people notice it while eating lunch, flossing before bed, or waking up with a strange rough spot on the side of a tooth. One minute the tooth feels normal, the next it catches the tongue, stings with cold water, or slices the inside of the cheek every time the jaw closes. It can feel minor at first, but a missing filling or newly sharp tooth edge deserves prompt attention, especially if pain, sensitivity, or swelling appear.

In practice, these problems fall into an awkward middle ground. They are not always dramatic enough for a hospital emergency room, but they are often urgent enough to justify calling an Emergency Dentist. The reason is simple. Once a filling comes out, the underlying tooth is exposed. That area may be softer, thinner, or already weakened by decay or a crack. A sharp edge can also signal that part of the tooth has fractured, and fractures do not reliably stay small.

The good news is that many of these cases are highly treatable when addressed early. A dentist can often smooth the edge, replace the filling, protect the tooth with a temporary or permanent restoration, or decide whether the problem points to a deeper crack or nerve issue. Waiting too long tends to narrow the options and increase cost, discomfort, and treatment time.

What a lost filling actually means

A filling does not simply fall out for no reason. Sometimes it has reached the end of its lifespan. Composite fillings, which are tooth colored, bond well and look natural, but they still wear over time, especially in areas that handle heavy chewing. Older silver amalgam fillings can last many years, but the tooth around them can weaken, or the filling can loosen as margins break down.

In other cases, the missing filling is a clue rather than the whole problem. There may be recurrent decay under the old restoration. The tooth may have cracked and released the filling. The bite may be too heavy on that spot, causing repeated stress. Patients who grind their teeth often show this pattern. I have seen people assume a piece of food came out, only to realize they were holding most of an old filling in their hand.

The sensation ***Vitality Dental Emergency Dentist*** afterward varies. Some teeth feel hollow or rough. Others become intensely sensitive to cold air, sweet foods, or pressure. If the filling was deep and close to the nerve, the change can be sharp and immediate. If a portion of the tooth broke with it, the edge may feel jagged enough to cut the tongue. That is one of the clearest signs that the tooth should be evaluated soon, even if the pain is not severe yet.

Why sharp tooth edges matter more than people expect

A sharp edge sounds like a comfort issue, but it often points to structural damage. Enamel is strong, but when it chips, the newly exposed area can be surprisingly abrasive. Patients commonly describe it as feeling like a tiny knife or a fishhook against the cheek. A small broken cusp on a back tooth may seem tolerable until chewing shifts pressure onto it and the crack travels deeper.

Soft tissue trauma is part of the problem. The tongue and cheeks do not leave rough areas alone. They constantly test them. Over a day or two, a minor chip can create a painful ulcer that makes speaking and eating miserable. The bigger concern is that a broken edge may indicate a tooth that cannot support normal function anymore. If that happens, a simple filling may no longer be enough. The tooth may need bonding, an onlay, a crown, or in some cases root canal treatment if the nerve has been compromised.

Front teeth deserve special attention because even a small chip changes the bite in ways people do not always notice. A patient may start avoiding contact on that side, clenching elsewhere, or using the chipped edge in a way that makes the fracture worse. Appearance matters too. Damage to a visible tooth often creates understandable stress, and emergency dental care can often restore shape quickly.

When it becomes urgent

Not every missing filling means a middle of the night crisis, but some signs move it into the urgent category. If you call an Emergency Dentist, the questions asked over the phone usually focus on pain, swelling, bleeding, trauma, and whether the tooth is broken in a way that affects eating or risks further damage.

Here are the situations that usually warrant same-day or next-day attention:

1. Strong pain that lingers, throbs, or wakes you at night
2. Visible swelling of the gum, face, or jaw
3. A sharp edge that is cutting the tongue or cheek repeatedly
4. A large portion of the tooth has broken away
5. Sensitivity or pain that makes it difficult to eat or drink

Those details matter because they help distinguish a repairable inconvenience from a progressing dental emergency. Swelling may indicate infection. Night pain often points to inflammation of the pulp, the tissue inside the tooth. A large missing section of tooth can fracture further if left unsupported. Even without dramatic pain, a back tooth that has lost a wide filling can collapse under normal chewing forces.

What to do before you are seen

There is a right way to manage the hours before a dental visit, and it is usually simple. The goal is to protect the tooth, reduce irritation, and avoid turning a small repair into a larger one.

If the filling came out whole, keep it if you can find it, although dentists do not usually reuse old fillings. It can still be helpful to see the size and shape of what was lost. Rinse your mouth gently with lukewarm water to clear debris. Avoid chewing on that side. If cold triggers pain, stick to foods and drinks that are closer to room temperature. Very sweet or sticky foods tend to aggravate exposed areas, so skip them for the moment.

For a sharp edge, orthodontic wax from a pharmacy can help cover the spot temporarily. Temporary dental repair material, also sold over the counter in many stores, can sometimes shield the area for a short period if used carefully. It is not a substitute for treatment, and it should not be packed aggressively into a deep hole, especially if that causes pain. If the tongue or cheek keeps rubbing a broken edge, a little wax can make a long wait far more tolerable.

A few practical precautions usually help:

1. Rinse gently after meals so food does not pack into the space
2. Use the opposite side of your mouth for chewing
3. Take an over the counter pain reliever if you normally can use it safely
4. Cover a cutting edge with dental wax if needed
5. Call a dentist promptly rather than waiting for the tooth to “settle down”

One point is worth stressing. Do not place aspirin directly on the gum or tooth. People still try this, usually because someone once suggested it as a home remedy. It can burn the tissue and does not solve the underlying problem.

What an Emergency Dentist looks for

At the appointment, the examination is usually quick but focused. The dentist will inspect the missing filling site, check whether the tooth is cracked, test the bite, and ask how long symptoms have been present. X-rays are Emergency Dentist often part of the visit because they show what cannot be seen from the surface, such as decay under an old restoration, a fracture pattern, or signs of infection near the root.

The distinction between a lost filling and a fractured tooth matters. If the remaining tooth structure is healthy and dry enough for bonding, the dentist may be able to replace the filling the same day. If the tooth is too weakened, rebuilding it with another standard filling may fail quickly. In those cases, a stronger restoration may be recommended.

Sharp edges are often smoothed right away if the fracture is minor and the tooth is otherwise stable. This can bring immediate relief. If the break is larger, smoothing alone may not be enough because reducing the edge further could remove too much structure or expose sensitive dentin. Then the goal shifts to restoring shape and support, sometimes temporarily until a full repair is completed.

Dentists also evaluate the pulp, the living tissue inside the tooth. A tooth that only reacts briefly to cold may still be calm enough for a straightforward restoration. A tooth that throbs spontaneously, remains painful after cold is removed, or hurts when tapped can be signaling more serious internal inflammation. When that happens, simply replacing the filling may not resolve the pain.

Common treatments and how dentists choose among them

The treatment depends on how much tooth remains, whether decay is present, and how the nerve is behaving. In a best-case scenario, the dentist cleans the area and places a new composite filling. This is common when the lost filling was small to medium in size and the tooth walls are still strong.

Bonding is often used when a front tooth chip or a modest broken edge needs reshaping. It can look excellent in skilled hands, and it is conservative because it preserves natural tooth. For back teeth, a larger defect may call for an onlay or crown. These options cover and reinforce weak cusps that might otherwise continue to crack.

Temporary restorations have a role, especially in emergency settings. If the tooth is too sensitive, bleeding, or difficult to isolate properly for a final bonded restoration, the dentist may place a temporary protective material and bring you back soon for the definitive repair. Patients sometimes underestimate how useful this is. A good temporary can buy comfort, stabilize the tooth, and prevent contamination while a longer-term plan is made.

If the nerve is irreversibly inflamed or infected, root canal treatment may be necessary before the final restoration. This surprises some people who thought they only needed “a new filling.” But if a lost filling exposed the tooth enough to trigger pulp damage, the emergency visit may be the moment that problem becomes obvious.

In severe cases where too little tooth remains, extraction enters the discussion. Dentists do not arrive at that recommendation lightly. It tends to come up when the fracture extends below the gumline, decay is extensive, or the remaining structure cannot support a predictable restoration. Catching the problem early often helps avoid that path.

The role of timing

A great many dental emergencies become more complicated because patients wait to see whether things improve on their own. Sometimes they do, briefly. Sensitivity can settle after the first day. The cheek may adapt around a rough edge. That lull can be misleading.

A missing filling creates a vulnerable site that traps food and plaque. Bacteria are given a shortcut into a prepared cavity that was once sealed. If the tooth already had a thin wall, every day of normal chewing increases the chance of a larger break. I have seen teeth that could have been refilled on Monday turn into crown cases by Friday, simply because one unsupported cusp gave way during dinner.

There is also the issue of pain patterns. Early nerve irritation can be intermittent. Once inflammation progresses, the pain tends to become less forgiving, often showing up at night or with temperature changes. Patients often describe this stage as a sudden worsening, but in truth the process was building quietly from the moment the restoration failed.

Special situations that change the plan

Not all mouths present the same way. Someone who grinds heavily at night may break fillings repeatedly unless the bite is adjusted and a night guard is considered. A patient with dry mouth, whether from medication or medical treatment, may have higher cavity risk around old fillings and need closer follow-up. Pregnancy can complicate timing and comfort, though urgent dental care is still appropriate when needed. Children and teens need a slightly different lens, especially if a chipped edge follows sports trauma or a fall.

Travel creates its own version of the problem. If a filling comes out before a long flight or during a trip, the temptation is to patch it and deal with it later. That can be reasonable for a day or two if symptoms are mild, but pain changes the equation quickly. Cabin pressure does not cause fillings to fall out, but teeth with inflamed

pulps can become more uncomfortable during air travel. When someone is far from their regular office, an Emergency Dentist can at least stabilize the tooth so the rest of the trip is manageable.

People with large older fillings deserve a careful discussion about expectations. When one of these restorations fails, the replacement is not always a simple swap. The old filling may have been holding together a tooth that has little natural structure left. In those cases, a crown is not overselling treatment. It is often the more durable and cost-effective choice over time.

What recovery usually feels like

After treatment, mild soreness is common, especially if the tooth was sensitive beforehand or the bite needed adjustment. A replaced filling should not feel high or interfere when you close your teeth together. If it does, call back. A restoration that hits first can make a tooth feel bruised within hours.

Sensitivity after a new filling can last a few days, sometimes a bit longer for deep cavities, but it should trend downward. Sharp pain when biting, or temperature pain that grows rather than fades, deserves re-evaluation. Most emergency dental repairs improve comfort quickly, but the timeline depends on what was actually wrong. A smoothed edge feels better almost immediately. A tooth that was close to needing root canal treatment may remain uncertain for a period, even after a careful repair.

Soft tissue ulcers from sharp edges usually heal once the source of trauma is removed. People are often surprised by how fast the tongue and cheeks recover. The mouth tends to heal well when it is no longer being poked all day.

Preventing the next emergency

Prevention is not glamorous, but it matters. Old fillings do not announce their retirement in advance. Regular exams help dentists spot marginal breakdown, cracks, and recurrent decay before a restoration suddenly fails. Bite forces matter more than many patients realize. If you clench or grind, protecting restored teeth can save you repeated emergency visits.

Diet and habits play a role too. Chewing ice, opening packages with teeth, and biting hard candy are classic ways to break both enamel and restorations. Sticky sweets can tug at compromised fillings. None of this means you need to eat nervously. It means weakened teeth benefit from a little respect.

There is also value in understanding your own dental history. If you know a particular tooth has a very large filling, previous crack lines, or recurring sensitivity, you will be quicker to act if something changes. Patients who treat that first rough spot or missing piece as important usually have simpler appointments and better outcomes.

A missing filling or sharp edge may begin as a nuisance, but teeth rarely become sturdier by being ignored. Quick assessment, sensible self-care, and timely treatment from an Emergency Dentist can turn a stressful moment into a straightforward repair. That is the real goal of emergency dental care, not just to stop the immediate discomfort, but to preserve as much healthy tooth as possible before a small break becomes a bigger problem.

Vitality Dental
1220 Coit Rd # 106, Plano, TX 75075, United States

+1 972-645-4100

FAQ About Emergency Dentist

What counts as a dental emergency?

A knocked-out tooth, significant mouth injury, severe tooth pain, or painful swelling warrants prompt evaluation. Call your dentist and describe your symptoms so the office can advise you on how urgently you need care.

What should I do if I knock out a permanent tooth?

Call a dentist immediately. Handle the tooth by the crown, avoid touching the root, and keep it moist. If you cannot put it back into the socket easily, transport it in milk or a tooth-preservation solution. Bring it with you to the appointment.

Should I go to the emergency room for a dental emergency?

Seek emergency medical care right away if you have trouble breathing or swallowing, uncontrolled bleeding, or a serious facial injury. For an isolated tooth injury or toothache without those symptoms, call a dentist promptly for guidance.